



Finitely-additive measures on the asymptotic foliations of a Markov compactum

Alexander I. Bufetov

► To cite this version:

Alexander I. Bufetov. Finitely-additive measures on the asymptotic foliations of a Markov compactum.
Moscow Mathematical Journal, 2014. hal-01256146

HAL Id: hal-01256146

<https://hal.science/hal-01256146>

Submitted on 14 Jan 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Finitely-additive measures on the asymptotic foliations of a Markov compactum.

Alexander I. Bufetov

1 Introduction.

1.1 Hölder cocycles over translation flows.

Let $\rho \geq 2$ be an integer, let M be a compact orientable surface of genus ρ , and let ω be a holomorphic one-form on M . Denote by $\mathfrak{m} = (\omega \wedge \bar{\omega})/2i$ the area form induced by ω and assume that $\mathfrak{m}(M) = 1$.

Let h_t^+ be the *vertical* flow on M (i.e., the flow corresponding to $\Re(\omega)$); let h_t^- be the *horizontal* flow on M (i.e., the flow corresponding to $\Im(\omega)$). The flows h_t^+ , h_t^- preserve the area \mathfrak{m} and are uniquely ergodic.

Take $x \in M$, $t_1, t_2 \in \mathbb{R}_+$ and assume that the closure of the set

$$\{h_{\tau_1}^+ h_{\tau_2}^- x, 0 \leq \tau_1 < t_1, 0 \leq \tau_2 < t_2\} \quad (1)$$

does not contain zeros of the form ω . Then the set (1) is called *an admissible rectangle* and denoted $\Pi(x, t_1, t_2)$. Let $\overline{\mathfrak{C}}$ be the semi-ring of admissible rectangles.

Consider the linear space \mathcal{Y}^+ of Hölder cocycles $\Phi^+(x, t)$ over the vertical flow h_t^+ which are invariant under horizontal holonomy. More precisely, a function $\Phi^+(x, t) : M \times \mathbb{R} \rightarrow \mathbb{C}$ belongs to the space \mathcal{Y}^+ if it satisfies:

1. $\Phi^+(x, t + s) = \Phi^+(x, t) + \Phi^+(h_t^+ x, s)$;
2. There exists $t_0 > 0$, $\theta > 0$ such that $|\Phi^+(x, t)| \leq t^\theta$ for all $x \in M$ and all $t \in \mathbb{R}$ satisfying $|t| < t_0$;
3. If $\Pi(x, t_1, t_2)$ is an admissible rectangle, then $\Phi^+(x, t_1) = \Phi^+(h_{t_2}^- x, t_1)$.

For example, if a cocycle Φ_1^+ is defined by $\Phi_1^+(x, t) = t$, then clearly $\Phi_1^+ \in \mathcal{Y}^+$.

In the same way define the space of \mathcal{Y}^- of Hölder cocycles $\Phi^-(x, t)$ over the horizontal flow h_t^- which are invariant under vertical holonomy, and set $\Phi_1^-(x, t) = t$.

Given $\Phi^+ \in \mathcal{Y}^+$, $\Phi^- \in \mathcal{Y}^-$, a finitely additive measure $\Phi^+ \times \Phi^-$ on the semi-ring $\overline{\mathfrak{C}}$ of admissible rectangles is introduced by the formula

$$\Phi^+ \times \Phi^-(\Pi(x, t_1, t_2)) = \Phi^+(x, t_1) \cdot \Phi^-(x, t_2). \quad (2)$$

In particular, for $\Phi^- \in \mathcal{Y}^-$, set $m_{\Phi^-} = \Phi_1^+ \times \Phi^-$:

$$m_{\Phi^-}(\Pi(x, t_1, t_2)) = t_1 \Phi^-(x, t_2). \quad (3)$$

For any $\Phi^- \in \mathcal{Y}^-$ the measure m_{Φ^-} satisfies $(h_t^+)_* m_{\Phi^-} = m_{\Phi^-}$ and is an invariant distribution in the sense of G. Forni [5], [6]. For instance, $m_{\Phi_1^-} = \mathbf{m}$.

A \mathbb{C} -linear pairing between \mathcal{Y}^+ and \mathcal{Y}^- is given, for $\Phi^+ \in \mathcal{Y}^+$, $\Phi^- \in \mathcal{Y}^-$, by the formula

$$\langle \Phi^+, \Phi^- \rangle = \Phi^+ \times \Phi^-(M) \quad (4)$$

The space of Lipschitz functions is not invariant under h_t^+ , and a larger function space $Lip_w^+(M, \omega)$ of weakly Lipschitz functions is introduced as follows. A bounded measurable function f belongs to $Lip_w^+(M, \omega)$ if there exists a constant C , depending only on f , such that for any admissible rectangle $\Pi(x, t_1, t_2)$ we have

$$\left| \int_0^{t_1} f(h_t^+ x) dt - \int_0^{t_1} f(h_t^+(h_{t_2}^- x)) dt \right| \leq C. \quad (5)$$

Let C_f be the infimum of all C satisfying (5). We norm $Lip_w^+(X)$ by setting

$$\|f\|_{Lip_w^+} = \sup_X f + C_f.$$

By definition, the space $Lip_w^+(M, \omega)$ contains all Lipschitz functions on M and is invariant under h_t^+ . We denote by $Lip_{w,0}^+(M, \omega)$ the subspace of $Lip_w^+(M, \omega)$ of functions whose integral with respect to \mathbf{m} is 0.

1.2 Flows along the stable foliation of a pseudo-Anosov diffeomorphism.

Assume that $\theta_1 > 0$ and a diffeomorphism $g : M \rightarrow M$ are such that

$$g^*(\Re(\omega)) = \exp(\theta_1)\Re(\omega); \quad g^*(\Im(\omega)) = \exp(-\theta_1)\Im(\omega). \quad (6)$$

The diffeomorphism g induces a linear automorphism g^* of the cohomology space $H^1(M, \mathbb{C})$. Denote by E^+ the expanding subspace of g^* (in other words, E^+ is the subspace spanned by vectors corresponding to Jordan cells of g^* with eigenvalues exceeding 1 in absolute value). The action of g on \mathcal{Y}^+ is given by $g^* \Phi^+(x, t) = \Phi^+(gx, \exp(\theta_1)t)$.

Proposition 1 *There exists a g^* -equivariant isomorphism between E^+ and \mathcal{Y}^+ .*

Theorem 1 *There exists a continuous mapping $\Xi^+ : Lip_w^+(M, \omega) \rightarrow \mathcal{Y}^+$ such that for any $f \in Lip_w^+(M, \omega)$, any $x \in X$ and any $T > 0$ we have*

$$\left| \int_0^T f \circ h_t^+(x) dt - \Xi^+(f)(x, T) \right| < C_\varepsilon \|f\|_{Lip_w^+} (1 + \log(1 + T))^{2\rho+1}.$$

The mapping Ξ^+ satisfies $\Xi^+(f \circ h_t^+) = \Xi^+(f)$ and $\Xi^+(f \circ g) = g^ \Xi^+(f)$.*

The mapping Ξ^+ is constructed as follows. By Proposition 1 applied to the flow h_t^- , there exists a g -equivariant isomorphism between \mathcal{Y}^- and the contracting space for the action of g^* on $H^1(M, \mathbb{C})$ (in other words, the subspace spanned by vectors corresponding to Jordan cells with eigenvalues strictly less than 1 in absolute value).

Proposition 2 *The pairing \langle, \rangle given by (4) is nondegenerate and g^* -invariant.*

Remark. Under the identification of \mathcal{Y}^+ and \mathcal{Y}^- with respective subspaces of $H^1(M, \mathbb{C})$, the pairing \langle, \rangle is taken to the cup-product on $H^1(M, \mathbb{C})$ (see Proposition 4.19 in Veech [14]).

If $f \in Lip_w^+(M, \omega)$, then f is Riemann-integrable with respect to m_{Φ^-} for any $\Phi^- \in \mathcal{Y}^-$ (see (30) for a precise definition of the integral). Assign to f a cocycle Φ_f^+ in such a way that for all $\Phi^- \in \mathcal{Y}^-$ we have

$$\langle \Phi_f^+, \Phi^- \rangle = \int_M f dm_{\Phi^-}. \quad (7)$$

By definition, $\Phi_{f \circ h_t^+}^+ = \Phi_f^+$. The mapping Ξ^+ of Theorem 1 is given by the formula

$$\Xi^+(f) = \Phi_f^+. \quad (8)$$

The first eigenvalue for the action of g^* on E^+ is $\exp(\theta_1)$ and is always simple. If its second eigenvalue has the form $\exp(\theta_2)$, where $\theta_2 > 0$, and is simple as well, then the following limit theorem holds for h_t^+ .

Given a bounded measurable function $f : X \rightarrow \mathbb{R}$ and $x \in X$, introduce a continuous function $\mathfrak{S}_n[f, x]$ on the unit interval by the formula

$$\mathfrak{S}_n[f, x](\tau) = \int_0^{\tau \exp(n\theta_1)} f \circ h_t^+(x) dt. \quad (9)$$

The functions $\mathfrak{S}_n[f, x]$ are $C[0, 1]$ -valued random variables on the probability space (M, \mathfrak{m}) .

Theorem 2 *If $g^*|_{E^+}$ has a simple, real second eigenvalue $\exp(\theta_2)$, $\theta_2 > 0$, then there exists a continuous functional $\alpha : Lip_w^+(M, \omega) \rightarrow \mathbb{R}$ and a compactly supported non-degenerate measure η on $C[0, 1]$ such that for any $f \in Lip_{w,0}^+(M, \omega)$ satisfying $\alpha(f) \neq 0$ the sequence of random variables*

$$\frac{\mathfrak{S}_n[f, x]}{\alpha(f) \exp(n\theta_2)}$$

converges in distribution to η as $n \rightarrow \infty$.

The functional α is constructed explicitly as follows. Under the assumptions of the theorem the action of g^* on E^- has a simple eigenvalue $\exp(-\theta_2)$; let $v(2)$ be the eigenvector with eigenvalue $\exp(-\theta_2)$, let $\Phi_2^- \in \mathcal{Y}^-$ correspond to $v(2)$ by Proposition 1 and $m_{\Phi_2^-}$ be given by (3); then

$$\alpha(f) = \int f dm_{\Phi_2^-}.$$

1.3 Generic translation flows.

Let $\rho \geq 2$ and let $\kappa = (\kappa_1, \dots, \kappa_\sigma)$ be a nonnegative integer vector such that $\kappa_1 + \dots + \kappa_\sigma = 2\rho - 2$. Denote by \mathcal{M}_κ the moduli space of Riemann surfaces of genus ρ endowed with a holomorphic differential of area 1 with singularities of orders k_1, \dots, k_σ (the *stratum* in the moduli space of holomorphic differentials), and let \mathcal{H} be a connected component of \mathcal{M}_κ . Denote by g_t the Teichmüller flow on \mathcal{H} (see [6], [8]), and let $\mathbb{A}(t, X)$ be the Kontsevich-Zorich cocycle over g_t [8].

Let \mathbb{P} be a g_t -invariant ergodic probability measure on \mathcal{H} . For $X \in \mathcal{H}$, $X = (M, \omega)$, let \mathcal{Y}_X^+ , \mathcal{Y}_X^- be the corresponding spaces of Hölder cocycles. Denote by E_X^+ the space spanned by the positive Lyapunov exponents of the Kontsevich-Zorich cocycle.

Proposition 3 *For \mathbb{P} -almost all $X \in \mathcal{H}$, we have $\dim \mathcal{Y}_X^+ = \dim \mathcal{Y}_X^- = \dim E_X^+$, and the pairing \langle, \rangle between \mathcal{Y}_X^+ and \mathcal{Y}_X^- is non-degenerate.*

Remark. In particular, if \mathbb{P} is the Masur-Veech “smooth” measure [10, 12], then $\dim \mathcal{Y}_X^+ = \dim \mathcal{Y}_X^- = \rho$.

Assign to $f \in Lip_w^+(M, \omega)$ a cocycle Φ_f^+ by (7).

Theorem 3 *For any $\varepsilon > 0$ there exists a constant C_ε depending only on \mathbb{P} such that for \mathbb{P} -almost every $X \in \mathcal{H}$, any $f \in Lip_w^+(X)$, any $x \in X$ and any $T > 0$ we have*

$$\left| \int_0^T f \circ h_t^+(x) dt - \Phi_f^+(x, T) \right| < C_\varepsilon \|f\|_{Lip_w^+} (1 + T^\varepsilon).$$

If both the first and the second Lyapunov exponent of the measure \mathbb{P} are positive and simple (as, by the Avila-Viana Theorem [2], is the case with the Masur-Veech “smooth” measure on \mathcal{H}), then the following limit theorem holds.

As before, consider a $C[0, 1]$ -valued random variable $\mathfrak{S}_t[f, x]$ on (M, \mathfrak{m}) defined by the formula

$$\mathfrak{S}_s[f, x](\tau) = \int_0^{\tau \exp(s)} f \circ h_t^+(x) dt.$$

Let $\|v\|$ be the Hodge norm in $H^1(M, \mathbb{R})$. Let $\theta_2 > 0$ be the second Lyapunov exponent of the Kontsevich-Zorich cocycle and let $v_2(X)$ be a Lyapunov vector corresponding to θ_2 (by our assumption, such a vector is unique up to scalar multiplication). Introduce a real-valued multiplicative cocycle $H_2(t, X)$ over g_t by the formula

$$H_2(t, X) = \frac{\|A(t, X)v_2(X)\|}{\|v_2(X)\|}. \quad (10)$$

Theorem 4 *Assume that both the first and the second Lyapunov exponent of the Kontsevich-Zorich cocycle with respect to the measure \mathbb{P} are positive and simple. Then for \mathbb{P} -almost any $X' \in \mathcal{H}$ there exists a non-degenerate compactly supported measure $\eta_{X'}$ on $C[0, 1]$ and, for \mathbb{P} -almost all $X, X' \in \mathcal{H}$, there exists a*

sequence of moments $s_n = s_n(X, X')$ such that the following holds. For \mathbb{P} -almost every $X \in \mathcal{H}$ there exists a continuous functional

$$\mathfrak{a}^{(X)} : Lip_w^+(X) \rightarrow \mathbb{R}$$

such that for \mathbb{P} -almost every X' and for any real-valued $f \in Lip_{w,0}^+(X)$ satisfying $\mathfrak{a}^{(X)}(f) \neq 0$, the sequence of $C[0,1]$ -valued random variables

$$\frac{\mathfrak{S}_{s_n}[f, x](\tau)}{(\mathfrak{a}^{(X)}(f))H_2(s_n, X)}$$

converges in distribution to $\eta_{X'}$ as $n \rightarrow \infty$.

Acknowledgements. W. A. Veech made the suggestion that G. Forni's invariant distributions for the vertical flow should admit a description in terms of cocycles for the horizontal flow, and I am greatly indebted to him. The observation that cocycles are dual objects to invariant distributions was made by G. Forni, and I am deeply grateful to him. I am deeply grateful to H. Nakada who pointed out the reference to S. Ito's work [7] to me. I am deeply grateful to J. Chaika, P. Hubert, Yu.S. Ilyashenko, E. Lanneau and R. Ryham for many helpful suggestions on improving the presentation. I am deeply grateful to A. Avila, X. Bressaud, B.M. Gurevich, A.V. Klimenko, V.I. Oseledets, Ya.G. Sinai, I.V. Vyugin, J.-C. Yoccoz for useful discussions. During the work on this paper, I was supported in part by the National Science Foundation under grant DMS 0604386 and by the Edgar Odell Lovett Fund at Rice University.

2 Asymptotic foliations of a Markov compactum.

2.1 Definitions and notation.

Let $m \in \mathbb{N}$ and let Γ be an oriented graph with m vertices $\{1, \dots, m\}$ and possibly multiple edges. We assume that for each vertex there is an edge starting from it and an edge ending in it.

Let $\mathcal{E}(\Gamma)$ be the set of edges of Γ . For $e \in \mathcal{E}(\Gamma)$ we denote by $I(e)$ its initial vertex and by $F(e)$ its terminal vertex. Let Q be the incidence matrix of Γ defined by the formula

$$Q_{ij} = \#\{e \in \mathcal{E}(\Gamma) : I(e) = i, F(e) = j\}.$$

By assumption, all entries of the matrix Q are positive. A finite word $e_1 \dots e_k$, $e_i \in \mathcal{E}(\Gamma)$, will be called *admissible* if $F(e_{i+1}) = I(e_i)$, $i = 1, \dots, k$.

To the graph Γ we assign a *Markov compactum* X_Γ , the space of bi-infinite paths along the edges:

$$X_\Gamma = \{x = \dots x_{-n} \dots x_0 \dots x_n \dots, x_n \in \mathcal{E}(\Gamma), F(x_{n+1}) = I(x_n)\}.$$

Remark. As Γ will be fixed throughout this section, we shall often omit the subscript Γ from notation and only insert it when the dependence on Γ is underlined.

Cylinders in X_Γ are subsets of the form $\{x : x_{n+1} = e_1, \dots, x_{n+k} = e_k\}$, where $n \in \mathbb{Z}$, $k \in \mathbb{N}$ and $e_1 \dots e_k$ is an admissible word. The family of all cylinders forms a semi-ring which we denote by \mathfrak{C} .

For $x \in X$, $n \in \mathbb{Z}$, introduce the sets

$$\gamma_n^+(x) = \{x' \in X_\Gamma : x'_t = x_t, t \geq n\}; \quad \gamma_n^-(x) = \{x' \in X_\Gamma : x'_t = x_t, t \leq n\};$$

$$\gamma_\infty^+(x) = \bigcup_{n \in \mathbb{Z}} \gamma_n^+(x); \quad \gamma_\infty^-(x) = \bigcup_{n \in \mathbb{Z}} \gamma_n^-(x).$$

The sets $\gamma_\infty^+(x)$ are leaves of the asymptotic foliation \mathcal{F}^+ on the space X_Γ ; the sets $\gamma_\infty^-(x)$ are leaves of the asymptotic foliation \mathcal{F}^- on X_Γ .

For $n \in \mathbb{Z}$ let \mathfrak{C}_n^+ be the collection of all subsets of X_Γ of the form $\gamma_n^+(x)$, $n \in \mathbb{Z}$, $x \in X$; similarly, \mathfrak{C}_n^- is the collection of all subsets of the form $\gamma_n^-(x)$. Set

$$\mathfrak{C}^+ = \bigcup_{n \in \mathbb{Z}} \mathfrak{C}_n^+; \quad \mathfrak{C}^- = \bigcup_{n \in \mathbb{Z}} \mathfrak{C}_n^-. \quad (11)$$

The collection \mathfrak{C}_n^+ is a semi-ring for any $n \in \mathbb{Z}$. Since every element of \mathfrak{C}_n^+ is a disjoint union of elements of \mathfrak{C}_{n+1}^+ , the collection \mathfrak{C}^+ is a semi-ring as well. The same statements hold for \mathfrak{C}_n^- and \mathfrak{C}^- .

Let $\exp(\theta_1)$ be the spectral radius of the matrix Q , and let $h = (h_1, \dots, h_m)$ be the unique positive eigenvector of Q : we thus have $Qh = \exp(\theta_1)h$. Let $\lambda = (\lambda_1, \dots, \lambda_m)$ be the positive eigenvector of the transpose matrix Q^t : we have $Q^t\lambda = \exp(\theta_1)\lambda$. The vectors λ, h are normalized as follows:

$$\sum_{i=1}^m \lambda_i = 1; \quad \sum_{i=1}^m \lambda_i h_i = 1. \quad (12)$$

Introduce a sigma-additive positive measure Φ_1^+ on the semi-ring \mathfrak{C}^+ by the formula

$$\Phi_1^+(\gamma_n^+(x)) = h_{F(x_n)} \exp((n-1)\theta_1) \quad (13)$$

and a sigma-additive positive measure Φ_1^- on the semi-ring \mathfrak{C}^- by the formula

$$\Phi_1^-(\gamma_n^-(x)) = \lambda_{I(x_n)} \exp(-n\theta_1). \quad (14)$$

Let $n \in \mathbb{Z}$, $k \in \mathbb{N}$, and let $e_1 \dots e_k$ be an admissible word. The Parry measure ν on X_Γ is defined by the formula

$$\nu(\{x : x_{n+1} = e_1, \dots, x_{n+k} = e_k\}) = \lambda_{I(e_k)} h_{F(e_1)} \exp(-k\theta_1). \quad (15)$$

The measures Φ_1^+, Φ_1^- are conditional measures of the Parry measure ν in the following sense. If $C \in \mathfrak{C}$, then $\gamma_\infty^+(x) \cap C \in \mathfrak{C}^+$, $\gamma_\infty^-(x) \cap C \in \mathfrak{C}^-$ for any $x \in C$, and we have

$$\nu(C) = \Phi_1^+(\gamma_\infty^+(x) \cap C) \cdot \Phi_1^-(\gamma_\infty^-(x) \cap C). \quad (16)$$

2.2 Finitely-additive measures on leaves of asymptotic foliations.

Given $v \in \mathbb{C}^m$, write

$$|v| = \sum_{i=1}^m |v_i|. \quad (17)$$

The norms of all matrices in this paper are understood with respect to this norm. Consider the direct-sum decomposition

$$\mathbb{C}^m = E^+ \oplus E^-,$$

where E^+ is spanned by Jordan cells of eigenvalues of Q with absolute value exceeding 1, and E^- is spanned by Jordan cells corresponding to eigenvalues of Q with absolute value at most 1. Let $v \in E^+$ and for all $n \in \mathbb{Z}$ set $v^{(n)} = Q^n v$ (note that $Q|_{E^+}$ is by definition invertible). Introduce a finitely-additive complex-valued measure Φ_v^+ on the semi-ring \mathfrak{C}^+ (defined in (11)) by the formula

$$\Phi_v^+(\gamma_{n+1}^+(x)) = (v^{(n)})_{F(x_{n+1})}. \quad (18)$$

The measure Φ_v^+ is invariant under holonomy along \mathcal{F}^- : by definition, we have the following

Proposition 4 *If $F(x_n) = F(x'_n)$, then $\Phi_v^+(\gamma_n^+(x)) = \Phi_v^+(\gamma_n^+(x'))$.*

The measures Φ_v^+ span a complex linear space, which we denote \mathcal{Y}^+ (or, sometimes, \mathcal{Y}_Γ^+ , when dependence on Γ is stressed.) The map

$$\mathcal{I} : v \rightarrow \Phi_v^+ \quad (19)$$

is an isomorphism between E^+ and \mathcal{Y}_Γ^+ .

For Q^t , we have the direct-sum decomposition

$$\mathbb{C}^m = \tilde{E}^+ \oplus \tilde{E}^-,$$

where \tilde{E}^+ is spanned by Jordan cells of eigenvalues of Q^t with absolute value exceeding 1, and \tilde{E}^- is spanned by Jordan cells corresponding to eigenvalues of Q^t with absolute value at most 1. As before, for $\tilde{v} \in \tilde{E}^+$ set $\tilde{v}^{(n)} = (Q^t)^n \tilde{v}$ for all $n \in \mathbb{Z}$, and introduce a finitely-additive complex-valued measure $\Phi_{\tilde{v}}^-$ on the semi-ring \mathfrak{C}^- (defined in (11)) by the formula

$$\Phi_{\tilde{v}}^-(\gamma_n^-(x)) = (\tilde{v}^{(-n)})_{I(x_n)}. \quad (20)$$

By definition, the measure $\Phi_{\tilde{v}}^-$ is invariant under holonomy along \mathcal{F}^+ : more precisely, we have the following

Proposition 5 *If $I(x_n) = I(x'_n)$, then $\Phi_{\tilde{v}}^-(\gamma_n^-(x)) = \Phi_{\tilde{v}}^-(\gamma_n^-(x'))$.*

Let \mathcal{Y}_Γ^- be the space spanned by the measures $\Phi_v^-, v \in \tilde{E}^+$. The map

$$\tilde{\mathcal{I}} : v \rightarrow \Phi_v^- \quad (21)$$

is an isomorphism between \tilde{E}^+ and \mathcal{Y}_Γ^- .

Let $\sigma : X_\Gamma \rightarrow X_\Gamma$ be the shift defined by $(\sigma x)_i = x_{i+1}$. The shift σ naturally acts on the spaces $\mathcal{Y}_\Gamma^+, \mathcal{Y}_\Gamma^-$: given $\Phi \in \mathcal{Y}_\Gamma^+$ (or \mathcal{Y}_Γ^-), the measure $\sigma_*\Phi$ is defined, for $\gamma \in \mathfrak{C}^+$, by the formula

$$\sigma_*\Phi(\gamma) = \Phi(\sigma\gamma).$$

From the definitions we obtain

Proposition 6 *The following diagrams are commutative:*

$$\begin{array}{ccc} E^+ & \xrightarrow{\mathcal{I}} & \mathcal{Y}_\Gamma^+ \\ \downarrow Q & & \uparrow \sigma^* \\ E^+ & \xrightarrow{\mathcal{I}} & \mathcal{Y}_\Gamma^+ \\ \\ \tilde{E}^+ & \xrightarrow{\tilde{\mathcal{I}}} & \mathcal{Y}_\Gamma^- \\ \downarrow Q^t & & \downarrow \sigma^* \\ \tilde{E}^+ & \xrightarrow{\tilde{\mathcal{I}}} & \mathcal{Y}_\Gamma^- \end{array}$$

2.3 Pairings.

Given $\Phi^+ \in \mathcal{Y}^+, \Phi^- \in \mathcal{Y}^-$, introduce, in analogy with (16), a finitely additive measure $\Phi^+ \times \Phi^-$ on the semi-ring \mathfrak{C} of cylinders in X_Γ : for any $C \in \mathfrak{C}$ and $x \in C$, set

$$\Phi^+ \times \Phi^-(C) = \Phi^+(\gamma_\infty^+(x) \cap C) \cdot \Phi^-(\gamma_\infty^-(x) \cap C). \quad (22)$$

Note that by Propositions 4, 5, the right-hand side in (22) does not depend on $x \in C$.

More explicitly, let $v \in E^+, \tilde{v} \in \tilde{E}^+, \Phi_v^+ = \mathcal{I}(v), \Phi_{\tilde{v}}^- = \tilde{\mathcal{I}}(\tilde{v})$. As above, denote $v^{(n)} = Q^n v, \tilde{v}^{(n)} = (Q^t)^n \tilde{v}$. Let $n \in \mathbb{Z}, k \in \mathbb{N}$ and let $e_1 \dots e_k$ be an admissible word. Then

$$\Phi_v^+ \times \Phi_{\tilde{v}}^-(\{x : x_{n+1} = e_1, \dots, x_{n+k} = e_k\}) = (v^{(n)})_{F(e_1)} (\tilde{v}^{(-n-k)})_{I(e_{n+k})}. \quad (23)$$

There is a natural \mathbb{C} -linear pairing \langle, \rangle between the spaces \mathcal{Y}_Γ^+ and \mathcal{Y}_Γ^- : for $\Phi^+ \in \mathcal{Y}_\Gamma^+, \Phi^- \in \mathcal{Y}_\Gamma^-$, set

$$\langle \Phi^+, \Phi^- \rangle = \Phi^+ \times \Phi^-(X_\Gamma). \quad (24)$$

From (23) we derive

Proposition 7 *Let $v \in E^+$, $\tilde{v} \in \tilde{E}^+$, $\Phi_v^+ = \mathcal{I}_\Gamma(v)$, $\Phi_{\tilde{v}}^- = \tilde{\mathcal{I}}_\Gamma(\tilde{v})$. Then*

$$\langle \Phi_v^+, \Phi_{\tilde{v}}^- \rangle = \sum_{i=1}^m v_i \tilde{v}_i. \quad (25)$$

In particular, the pairing \langle, \rangle is non-degenerate and σ^ -invariant.*

In particular, for $\Phi^- \in \mathcal{Y}^-$ denote

$$m_{\Phi^-} = \Phi_1^+ \times \Phi^-. \quad (26)$$

2.4 Weakly Lipschitz Functions.

Introduce a function space $Lip_w^+(X)$ in the following way. A bounded Borel-measurable function $f : X \rightarrow \mathbb{C}$ belongs to the space $Lip_w^+(X)$ if there exists a constant $C > 0$ such that for all $n \geq 0$ and any $x, x' \in X$ satisfying $F(x_{n+1}) = F(x'_{n+1})$, we have

$$\left| \int_{\gamma_n^+(x)} f d\Phi_1^+ - \int_{\gamma_n^+(x')} f d\Phi_1^+ \right| \leq C. \quad (27)$$

If C_f be the infimum of all C satisfying (27), then we norm $Lip_w^+(X)$ by setting

$$\|f\|_{Lip_w^+} = \sup_X f + C_f.$$

As before, let $Lip_{w,0}^+(X)$ be the subspace of $Lip_w^+(X)$ of functions whose integral with respect to ν is zero.

Take $\Phi^- \in \mathcal{Y}^-$. Any function $f \in Lip_w^+(X)$ is integrable with respect to the measure m_{Φ^-} , defined by (26), in the following sense. Let $\tilde{v} \in E^-$ be the vector corresponding to Φ^- by (20) and let $\tilde{v}^{(n)} = (Q^t)^n \tilde{v}$. Recall that

$$|\tilde{v}^{(-n)}| \rightarrow 0 \text{ exponentially fast as } n \rightarrow \infty. \quad (28)$$

Take arbitrary points $x_i^{(n)} \in X$, $n \in \mathbb{N}$ satisfying

$$F((x_i^{(n)})_n) = i, \quad i = 1, \dots, m. \quad (29)$$

and consider the expression

$$\sum_{i=1}^m \left(\int_{\gamma_n^+(x_i^{(n)})} f d\Phi_1^+ \right) \cdot (\tilde{v}^{(1-n)})_i. \quad (30)$$

By (27) and (28), as $n \rightarrow \infty$ the expression (30) tends to a limit which does not depend on the particular choice of $x_i^{(n)}$ satisfying (29). This limit is denoted

$$m_{\Phi^-}(f) = \int_X f dm_{\Phi^-}.$$

Introduce a measure $\Phi_f^+ \in \mathcal{Y}^+$ by requiring that for any $\Phi^- \in \mathcal{Y}^-$ we have

$$\langle \Phi_f^+, \Phi^- \rangle = \int_X f dm_{\Phi^-}. \quad (31)$$

Note that the mapping $\Xi^+ : Lip_w^+(X) \rightarrow \mathcal{Y}^+$ given by $\Xi^+(f) = \Phi_f^+$ is continuous by definition and satisfies

$$\Xi^+(f \circ \sigma) = \sigma^* \Xi^+(f). \quad (32)$$

From the definitions we also have

Proposition 8 *Let $\Phi^+(1), \dots, \Phi^+(r)$ be a basis in \mathcal{Y}^+ and let $\Phi^-(1), \dots, \Phi^-(r)$ be the dual basis in \mathcal{Y}^- with respect to the pairing \langle, \rangle . Then for any $f \in Lip_w^+(X)$ we have*

$$\Phi_f^+ = \sum_{i=1}^r (m_{\Phi^-(i)}(f)) \Phi^+(i).$$

2.5 Approximation.

Let Θ be a finitely-additive complex-valued measure on the semi-ring \mathfrak{C}_0^+ . Assume that there exists a constant $\delta(\Theta)$ such that for all $x, x' \in X$ and all $n \geq 0$ we have

$$|\Theta(\gamma_n^+(x)) - \Theta(\gamma_n^+(x'))| \leq \delta(\Theta) \text{ if } F(x_{n+1}) = F(x'_{n+1}). \quad (33)$$

In this case Θ will be called a *weakly Lipschitz measure*.

Lemma 1 *There exists a constant C_Γ depending only on Γ such that the following is true. Let Θ be a weakly Lipschitz finitely-additive complex-valued measure on the semi-ring \mathfrak{C}_0^+ . Then there exists a unique $\Phi^+ \in \mathcal{Y}_\Gamma^+$ such that for all $x \in X$ and all $n > 0$ we have*

$$|\Theta(\gamma_n^+(x)) - \Phi^+(\gamma_n^+(x))| \leq C_\Gamma \delta(\Theta) n^{m+1}. \quad (34)$$

Assign to the graph Γ the Markov compactum Y_Γ of one-sided infinite sequences of edges:

$$Y = \{y = y_1 \dots y_n \dots : y_n \in \mathcal{E}(\Gamma), F(y_{n+1}) = I(y_n)\},$$

and, as before, let σ be the shift on Y_Γ : $(\sigma y)_i = y_{i+1}$. For $y, y' \in Y_\Gamma$, write $y' \searrow y$ if $\sigma y' = y$.

Lemma 1 will be derived from

Lemma 2 *There exists a constant C_Γ depending only on Γ such that the following is true. Let φ_n be a sequence of measurable complex-valued functions on Y_Γ . Assume that there exists a constant δ such that for all $y \in Y$ and all $n \geq 0$ we have*

$$|\varphi_{n+1}(y) - \sum_{y' \searrow y} \varphi_n(y')| \leq \delta \quad (35)$$

and for all $n \geq 0$ and all $y, \tilde{y} \in Y_\Gamma$ satisfying $F(y_1) = F(\tilde{y}_1)$ we have

$$|\varphi_n(y) - \varphi_n(\tilde{y})| \leq \delta. \quad (36)$$

Then there exists a unique $v \in E^+$ such that for all $y \in Y$ and all $n > 0$ we have

$$|\varphi_n(y) - (Q^n v)_{F(y_{n+1})}| \leq C_\Gamma \delta n^{m+1}. \quad (37)$$

Proof of Lemma 2. Take arbitrary points $y(i) \in Y_\Gamma$ in such a way that

$$F(y(i)_1) = i.$$

Introduce a sequence of vectors $v(n) \in \mathbb{C}^m$ by the formula

$$v(n)_i = \varphi_n(y(i)).$$

From (36) for any $y \in Y$ we have

$$|\varphi_n(y) - v(n)_{F(y_1)}| \leq \delta,$$

and from (35), (36) we have

$$|Qv(n) - v(n+1)| \leq \delta \cdot \|Q\|.$$

To prove Lemma 2, it suffices now to establish the following

Proposition 9 *Let V be a finite-dimensional complex linear space, let $S : V \rightarrow V$ be a linear operator and let $V^+ \subset V$ be the subspace spanned by vectors corresponding to Jordan cells of S with eigenvalues exceeding 1 in absolute value. There exists a constant $C > 0$ depending only on S such that the following is true. Assume that the vectors $v(n) \in V$, $n \in \mathbb{N}$, satisfy*

$$|Sv(n) - v(n+1)| < \delta$$

for all $n \in \mathbb{N}$ and some constant $\delta > 0$. Then there exists a unique $v \in V^+$ such that for all $n \in \mathbb{N}$ we have

$$|S^n v - v(n)| \leq C \cdot \delta \cdot n^{\dim V - \dim V^+ + 1}. \quad (38)$$

Proof of Proposition 9. By definition, the subspace V^+ is S -invariant and S is invertible on V^+ ; we have furthermore that $|Q^{-n}v| \rightarrow 0$ exponentially fast as $n \rightarrow \infty$. Let V^- be the subspace spanned by Jordan cells corresponding to eigenvalues of absolute value at most 1; for $v \in V^-$, we have $|Q^n v| < C n^{\dim V - \dim V^+}$ as $n \rightarrow \infty$. We have the decomposition $V = V^+ \oplus V^-$. Let

$$u(0) = v(0), u(n+1) = v(n+1) - Sv(n).$$

Decompose $u(n) = u^+(n) + u^-(n)$, where $u^+(n) \in V^+$, $u^-(n) \in V^-$. Denote

$$v^+(n+1) = u^+(n+1) + Su^+(n) + \cdots + S^n u^+(1);$$

$$v^-(n+1) = u^-(n+1) + Su^-(n) + \dots + S^n u^-(1);$$

$$v = u^+(0) + S^{-1}u^+(1) + \dots + S^{-n}u^+(n) + \dots$$

By definition, $|v^-(n+1)|$ is bounded above by $C\delta n^{\dim V - \dim V^+ + 1}$ and there exists \tilde{C} such that $|S^n v - v^+(n)| < \tilde{C}\delta$ for all $n \in \mathbb{N}$, whence (38) follows. Uniqueness of v follows from the fact that for any nonzero $v' \in V^+$ the sequence $|S^n v'|$ grows exponentially as $n \rightarrow \infty$. Proposition 9 and Lemmas 1, 2 are proved completely.

Let $f \in Lip_w^+(X)$. We then have a measure Θ_f on the semi-ring \mathfrak{C}_0^+ given, for $\gamma \in \mathfrak{C}_0^+$, by the formula

$$\Theta_f(\gamma) = \int_{\gamma} f d\Phi_1^+.$$

By (27), the measure Θ_f satisfies the assumptions of Lemma 1. Let $\Xi_f^+ \in \mathcal{Y}^+$ be the measure assigned to Θ_f by Lemma 1.

Lemma 3 *Let $f \in Lip_w^+(X)$, $\Phi^- \in \mathcal{Y}_\Gamma^-$. Then*

$$\langle \Xi_f^+, \Phi^- \rangle = \int_X f dm_{\Phi^-}. \quad (39)$$

Proof: Choose the points $x_i^{(n)} \in X$ satisfying (29). As above, let $\tilde{v} \in E^-$ be the vector corresponding to Φ^- by (20) and let $\tilde{v}^{(n)} = (Q^t)^n \tilde{v}$, $n \in \mathbb{Z}$. For any $\varepsilon > 0$ and $n > 0$ sufficiently large, by definition, we have

$$|m_{\Phi^-}(f) - \sum_{i=1}^m \left(\int_{\gamma_n^+(x_i^{(n)})} f d\Phi_1^+ \right) \cdot (\tilde{v}^{(-n)})_i| < \varepsilon. \quad (40)$$

By definition of Ξ_f^+ and Lemma 1 we have

$$\left| \sum_{i=1}^m \left(\int_{\gamma_n^+(x_i^{(n)})} f d\Phi_1^+ \right) \cdot (\tilde{v}^{(-n)})_i - \sum_{i=1}^m (\Xi_f^+(\gamma_n^+(x_i^{(n)})) \cdot (\tilde{v}^{(-n)})_i \right| < C_\Gamma \cdot n^{m+1} |\tilde{v}_i^{(-n)}|,$$

and, by (28), the right-hand side tends to 0 exponentially fast as $n \rightarrow \infty$.

It remains to notice that, by definition,

$$\sum_{i=1}^m (\Xi_f^+(\gamma_n^+(x_i^{(n)})) \cdot (\tilde{v}^{(-n)})_i = \langle \Xi_f^+, \Phi^- \rangle,$$

and the Lemma is proved completely.

We have thus established that $\Xi_f^+ = \Phi_f^+$, where Φ_f^+ is given by (31).

2.6 Orderings.

Following S. Ito [7], A.M. Vershik [15, 16], assume that a partial order \mathfrak{o} is given on $\mathcal{E}(\Gamma)$ in such a way that edges starting at a given vertex are ordered linearly, while edges starting at different vertices are not comparable. An edge will be called *maximal* (with respect to \mathfrak{o}) if there does not exist a greater edge; *minimal*, if there does not exist a smaller edge; and an edge e will be called *the successor* of e' if $e > e'$ but there does not exist e'' such that $e > e'' > e'$.

The ordering \mathfrak{o} is extended to a partial ordering of X_Γ : we write $x < x'$ if there exists $l \in \mathbb{Z}$ such that $x_l < x'_l$ and $x_n = x'_n$ for all $n > l$. Under this ordering each leaf γ_∞^+ of the foliation \mathcal{F}^+ is linearly ordered, while points lying on different leaves are not comparable.

Let $Max(\mathfrak{o})$ be the set of points $x \in X$, $x = (x_n)_{n \in \mathbb{Z}}$, such that each x_n is a maximal edge. Similarly, $Min(\mathfrak{o})$ denotes the set of points $x \in X$, $x = (x_n)_{n \in \mathbb{Z}}$, such that each x_n is a minimal edge. Since edges starting at a given vertex are ordered linearly, the cardinalities of $Max(\mathfrak{o})$ and $Min(\mathfrak{o})$ do not exceed m .

If a leaf γ_∞^+ does not intersect $Max(\mathfrak{o})$, then it does not have a maximal element; similarly, if γ_∞^+ does not intersect $Min(\mathfrak{o})$, then it does not have a minimal element.

For $x(1), x(2) \in \gamma_\infty^+$, let

$$[x(1), x(2)] = \{x' \in \gamma_\infty^+ : x(1) \leq x' \leq x(2)\}.$$

The sets $(x(1), x(2))$, $[x(1), x(2))$, $(x(1), x(2)]$ are defined similarly.

Proposition 10 *Let $x \in X$. If $\gamma_\infty^+(x) \cap Max(\mathfrak{o}) = \emptyset$, then for any $t \geq 0$ there exists a point $x' \in \gamma_\infty^+(x)$ such that*

$$\Phi_1^+([x, x']) = t. \quad (41)$$

Proof. Let $V(x) = \{t : \exists x' \geq x : \Phi_1^+([x, x']) = t\}$. Since $\gamma_\infty^+(x) \cap Max(\mathfrak{o}) = \emptyset$, for any n there exists $x'' \in \gamma_\infty^+(x)$ such that all points in $\gamma_n^+(x'')$ are greater than x . Since $\Phi_1^+(\gamma_n^+(x''))$ grows exponentially, uniformly in x'' , as $n \rightarrow \infty$, the set $V(x)$ is unbounded. Furthermore, since $\Phi_1^+(\gamma_n^+(x''))$ decays exponentially, uniformly in x'' , as $n \rightarrow -\infty$, the set $V(x)$ is dense in \mathbb{R}_+ . Finally, by compactness of X , the set $V(x)$ is closed, which concludes the proof of the Proposition.

A similar proposition, proved in the same way, holds for negative t .

Proposition 11 *Let $x \in X$. If $\gamma_\infty^+(x) \cap Min(\mathfrak{o}) = \emptyset$, then for any $t \geq 0$ there exists a point $x' \in \gamma_\infty^+(x)$ such that*

$$\Phi_1^+([x', x]) = t. \quad (42)$$

Define an equivalence relation \sim on X by writing $x \sim x'$ if $x \in \gamma_\infty^+(x')$ and $\Phi_1^+([x, x']) = \Phi_1^+([x', x]) = 0$. The equivalence classes admit the following explicit description, which is clear from the definitions.

Proposition 12 *Let $x, x' \in X$ be such that $x \in \gamma_\infty^+(x')$, $x < x'$ and $\Phi_1^+([x, x']) = 0$. Then there exists $n \in \mathbb{Z}$ such that*

1. x'_n is a successor of x_n ;
2. x is the maximal element in $\gamma_n(x)$;
3. x' is the minimal element in $\gamma_n(x')$.

In other words, $\Phi_1^+([x, x']) = 0$ if and only if $(x, x') = \emptyset$. In particular, equivalence classes consist at most of two points and, ν -almost surely, of only one point.

Denote $X_\circ = X/\sim$, let $\pi_\circ : X \rightarrow X_\circ$ be the projection map and set $\nu_\circ = (\pi_\circ)_*\nu$. The probability spaces (X_\circ, ν_\circ) and (X, ν) are measurably isomorphic; in what follows, we shall often omit the index \circ . The foliations \mathcal{F}^+ and \mathcal{F}^- descend to the space X_\circ ; we shall denote their images on X_\circ by the same letters and, as before, denote by $\gamma_\infty^+(x)$, $\gamma_\infty^-(x)$ the leaves containing $x \in X_\circ$.

Now let $x \in X_\circ$ satisfy $\gamma_\infty^+(x) \cap \text{Max}(\circ) = \emptyset$. By Proposition 10, for any $t \geq 0$ there exists a unique x' satisfying (41). Denote $h_t^+(x) = x'$. Similarly, if $x \in X_\circ$ satisfy $\gamma_\infty^-(x) \cap \text{Min}(\circ) = \emptyset$. By Proposition 11, for any $t \geq 0$ there exists a unique x' satisfying (42). Denote $h_{-t}^-(x) = x'$.

We thus obtain a flow h_t^+ , which is well-defined on the set

$$X_\circ \setminus \left(\bigcup_{x \in \text{Max}(\circ) \cup \text{Min}(\circ)} \gamma_\infty^+(x) \right),$$

and, in particular, ν -almost surely on X_\circ . By (16), the flow h_t^+ preserves the measure ν .

More generally, it is clear from the definitions that for any $\Phi^- \in \mathcal{Y}^-$, the measure m_{Φ^-} , defined by (26), satisfies

$$(h_t^+)_* m_{\Phi^-} = m_{\Phi^-},$$

similarly to G. Forni's invariant distributions [5], [6].

Remark. S.Ito in [7] gives a construction of a flow similar to the one above. The flow h_t^+ is a continuous-time analogue of a Vershik automorphism [15] (of which a variant also occurs in Ito's work [7]), and, in fact, is a suspension flow over the corresponding Vershik's automorphism, a point of view adopted in [4].

2.7 Decomposition of Arcs.

We assume that an ordering \circ is fixed on Γ . Denote by $\mathfrak{C}(\circ)$ the semi-ring of subsets of X_Γ of the form $[x, x')$, where $x < x'$. Any measure $\Phi^+ \in \mathcal{Y}^+$ can be extended to $\mathfrak{C}(\circ)$ in the following way.

Let \mathfrak{R}_n^+ be the ring generated by the semi-ring \mathfrak{C}_n^+ . For $\gamma \in \mathfrak{C}(\circ)$, denote by $\gamma(n)$ the smallest (by inclusion) element of the ring \mathfrak{R}_{-n}^+ containing γ and

let $\hat{\gamma}(n)$ be the greatest (by inclusion) element of the ring \mathfrak{R}_{-n}^+ contained in γ (possibly, $\hat{\gamma}(n) = \emptyset$). By definition,

$$\hat{\gamma}(n) \subset \hat{\gamma}(n+1) \subset \gamma(n+1) \subset \gamma(n);$$

$$\gamma(n) \setminus \hat{\gamma}(n) = \bigsqcup_{i=1}^{l_n} \gamma_i^{(n)}, \quad (43)$$

where $\gamma_i^{(n)} \in \mathfrak{C}_{-n}^+$, $l_n \leq \|Q\|$, and

$$\gamma(n) \setminus \gamma(n+1) = \bigsqcup_{i=1}^{L_n} \gamma_i^{(n+1)}, \quad (44)$$

where $\gamma_i^{(n+1)} \in \mathfrak{C}_{-n-1}^+$, $L_n \leq 2\|Q\|$.

By definition, if $\Phi^+ \in \mathcal{Y}^+$, then there are only m possible values of $\Phi^+(\gamma)$ for $\gamma \in \mathfrak{C}_{-n}^+$, and the maximum of these decays exponentially as $n \rightarrow \infty$. We thus have

Proposition 13 *There exists positive constants C_Γ , depending only on Γ , such that the following is true. Let $v_0 = 0$, $v_1, \dots, v_l \in E^+$, $Qv_i = \exp(\theta)v_i + v_{i-1}$. Assume $v \in \mathbb{C}v_1 \oplus \dots \oplus \mathbb{C}v_l$ satisfies $|v| = 1$ and let $\Phi_v^+ = \mathcal{I}_\Gamma(v)$. Then for any $\gamma \in \mathfrak{C}(\mathfrak{o})$ we have*

$$|\Phi_v^+(\gamma(n)) - \Phi_v^+(\gamma(n+1))| \leq C_\Gamma n^{l-1} \exp(-(\Re\theta)n);$$

$$|\Phi_v^+(\hat{\gamma}(n)) - \Phi_v^+(\hat{\gamma}(n+1))| \leq C_\Gamma n^{l-1} \exp(-(\Re\theta)n).$$

decay exponentially as $n \rightarrow \infty$. In particular, if $v \in E^+$, $Qv = \exp(\theta)v$, $|v| = 1$, then

$$|\Phi_v^+(\gamma(n)) - \Phi_v^+(\gamma(n+1))| \leq C_\Gamma \exp(-(\Re\theta)n);$$

$$|\Phi_v^+(\hat{\gamma}(n)) - \Phi_v^+(\hat{\gamma}(n+1))| \leq C_\Gamma \exp(-(\Re\theta)n).$$

Consequently, for any $\Phi^+ \in \mathcal{Y}^+$, $\gamma \in \mathfrak{C}(\mathfrak{o})$, the sequence $\Phi^+(\gamma(n))$ converges as $n \rightarrow \infty$, and we set

$$\Phi^+(\gamma) = \lim_{n \rightarrow \infty} \Phi^+(\gamma(n)).$$

By (43), we also have

$$\Phi^+(\gamma) = \lim_{n \rightarrow \infty} \Phi^+(\hat{\gamma}(n)).$$

Proposition 14 *The measure Φ^+ is finitely-additive on $\mathfrak{C}(\mathfrak{o})$.*

Proof: Let $v \in E^+$ be such that $\Phi^+ = \Phi_v^+$ and let $\gamma_0, \gamma_1, \dots, \gamma_k \in \mathfrak{C}(\mathfrak{o})$ satisfy

$$\gamma_0 = \bigsqcup_{i=1}^k \gamma_i.$$

Consider the arcs $\gamma_0(n), \gamma_1(n), \dots, \gamma_k(n)$. We have

$$\gamma_0(n) \subset \bigcup_{i=1}^k \gamma_i(n). \quad (45)$$

and decompose

$$\gamma_i(n) = \bigsqcup \gamma_{ij}(n+1),$$

where $\gamma_{ij}(n+1) \in \mathfrak{C}_{-n-1}^+$.

By (45), each of the arcs $\gamma_{0j}(n+1)$ is also encountered among the arcs $\gamma_{ij}(n+1)$ (possibly, more than once, but not more than k times). Consider the collection $\gamma_{ij}(n+1)$ and cross out all the arcs $\gamma_{0j}(n+1)$; by maximality, and since our ordering is linear on each leaf of the foliation \mathcal{F}^+ , there will remain not more than $2k\|Q\|$ arcs, whence we obtain

$$\left| \sum_{i=1}^k \Phi^+(\gamma_i(n)) - \Phi^+(\gamma_0(n)) \right| \leq 2k\|Q\| \cdot |Q^{-n-1}v|,$$

and, since the right-hand side decays exponentially as $n \rightarrow \infty$, the Proposition is proved.

Lemma 4 *There exists a constant C_Γ depending only on Γ such that the following is true. Let $f \in Lip_w^+(X_\Gamma)$ and let $\Phi_f^+ \in \mathcal{Y}^+$ be given by (31). For any $\gamma \in \mathfrak{C}(\mathfrak{o})$ we have*

$$\left| \int_\gamma f d\Phi_1^+ - \Phi_f^+(\gamma) \right| \leq C_\Gamma \|f\|_{Lip_w^+} (1 + \log(1 + \Phi_1^+(\gamma)))^{m+1}. \quad (46)$$

Indeed, for $\gamma \in \mathfrak{C}^+$ this follows from Lemma 1, and for all other arcs from Proposition 13.

2.8 Ergodic averages of the flow h_t^+ .

Let $\Phi^+ \in \mathcal{Y}^+$ and denote $\Phi^+[x, t] = \Phi_t^+([x, h_t^+x])$. The function $\Phi^+(x, t)$ is an additive cocycle over the flow h_t^+ . Let $f \in Lip_w^+(X_\Gamma)$, and let Φ_f^+ be defined by (31). By definition, $\Phi_{f \circ h_t^+} = \Phi_f^+$; recall from (32) that $\Phi_{f \circ \sigma} = \sigma^* \Phi_f^+$. Lemma 4 implies

Theorem 5 *There exists a positive constant C_Γ depending only on Γ such that for any $f \in Lip_w^+(X_\Gamma)$, for all $x \in X$ and all $T > 0$ we have*

$$\left| \int_0^T f \circ h_t^+(x) dt - \Phi_f^+(x, T) \right| \leq C_\Gamma \|f\|_{Lip} (1 + \log(1 + T))^{m+1}.$$

Given a bounded measurable function $f : X \rightarrow \mathbb{R}$ and $x \in X$, introduce a continuous function $\mathfrak{S}_n[f, x]$ on the unit interval by the formula

$$\mathfrak{S}_n[f, x](\tau) = \int_0^{\tau \exp(n\theta_1)} f \circ h_t^+(x) dt. \quad (47)$$

The functions $\mathfrak{S}_n[f, x]$ are $C[0, 1]$ -valued random variable on the probability space (X_Γ, ν_Γ) .

Theorem 6 *If Q has a simple real second eigenvalue $\exp(\theta_2)$, $\theta_2 > 0$, then there exists a continuous functional $\alpha : Lip_w^+(X) \rightarrow \mathbb{R}$ and a compactly supported non-degenerate measure η on $C[0, 1]$ such that for any $f \in Lip_{w,0}^+(X)$ satisfying $\alpha(f) \neq 0$ the sequence of random variables*

$$\frac{\mathfrak{S}_n[f, x]}{\alpha(f) \exp(n\theta_2)}$$

converges in distribution to η as $n \rightarrow \infty$.

Remark. Compactness of the support of η is understood in the sense of the Tchebycheff topology on $C[0, 1]$. Nondegeneracy of the measure η means that if $\varphi \in C[0, 1]$ is distributed according to η , then for any $t_0 \in (0, 1]$ the distribution of the real-valued random variable $\varphi(t_0)$ is not concentrated at a single point.

The measure η is constructed as follows: let v_2 be an eigenvector with eigenvalue $\exp(\theta_2)$, set $\Phi_2^+ = \mathcal{I}(v_2)$ (see (19)); then η is the distribution of $\Phi_2^+(x, \tau)$, $0 \leq \tau \leq 1$, considered as a $C[0, 1]$ -valued random variable on the space X_Γ, ν_Γ . The functional $\alpha(f)$ is constructed as follows: under the assumptions of Theorem 6, the matrix Q^t also has the simple real second eigenvalue $\exp(\theta_2)$; let \tilde{v}_2 be the eigenvector with eigenvalue $\exp(\theta_2)$, normalized in such a way that $\sum_{i=1}^m (v_2)_i (\tilde{v}_2)_i = 1$; set $\Phi_2^- = \tilde{\mathcal{I}}(\tilde{v}_2)$ (see (21)), and let $m_{\Phi_2^-}$ be given by (26); then

$$\alpha(f) = \int f dm_{\Phi_2^-}.$$

2.9 The diagonalizable case.

As an illustration, consider the case when $Q|_{E^+}$ is diagonalizable with eigenvalues $\exp(\theta_i)$, $i = 1, \dots, r$, $\Re(\theta_i) > 0$. The Perron-Frobenius vector h corresponds to $\exp(\theta_1)$; let v_2, \dots, v_r be eigenvectors corresponding to $\exp(\theta_i)$: thus $Qv_i = \exp(\theta_i)v_i$, $i = 2, \dots, r$ and

$$E^+ = \mathbb{C}h \oplus \mathbb{C}v_2 \oplus \dots \oplus \mathbb{C}v_r.$$

We have a similar direct-sum representation for Q^t :

$$\tilde{E}^+ = \mathbb{C}\lambda \oplus \mathbb{C}\tilde{v}_2 \oplus \dots \oplus \mathbb{C}\tilde{v}_r,$$

where $Q^t \tilde{v}_i = \exp(\theta_i) \tilde{v}_i$, $i = 2, \dots, r$. For $i \neq j$ we have

$$\sum_{l=1}^m (v_i)_l (\tilde{v}_j)_l = 0, \quad (48)$$

and, for normalization, let us assume that for all $i = 1, \dots, r$ we have

$$\sum_{l=1}^m (v_i)_l (\tilde{v}_i)_l = 1. \quad (49)$$

Let $\Phi_i^+ = \mathcal{I}(v_i)$, $\Phi_i^- = \tilde{\mathcal{I}}(\tilde{v}_i)$, $i = 2, \dots, r$. Since $\Phi_1^+ = \mathcal{I}(h)$, the measures Φ_i^+ , $i = 1, \dots, r$, form a basis in \mathcal{Y}^+ , for which the measures $\Phi_1^- = \tilde{\mathcal{I}}(\lambda)$, $\Phi_2^-, \dots, \Phi_r^-$ form a dual basis in \mathcal{Y}^- .

For $i = 1, \dots, r$, from (26) we have the measures $m_{\Phi_i^-} = \Phi_1^+ \times \Phi_i^-$. For instance, $m_{\Phi_1^-} = \nu$. Theorem 5 now implies

Corollary 1 *For any $f \in Lip_w^+(X_\Gamma)$ we have*

$$\left| \int_0^T f \circ h_t^+(x) dt - T \int_X f d\nu - \sum_{i=2}^r \Phi_i^+(x, T) (m_{\Phi_i^-}(f)) \right| \leq C_\Gamma \|f\|_{Lip} (1 + \log(1+T))^{m+1},$$

where C_Γ is a constant depending only on Γ .

For the action of the shift we have:

$$(\sigma)_* \Phi_i^+ = \exp(-\theta_i) \Phi_i^+, \quad i = 1, \dots, r; \quad (50)$$

$$(\sigma)_* \Phi_i^- = \exp(\theta_i) \Phi_i^-, \quad i = 1, \dots, r. \quad (51)$$

Corollary 1 now yields

$$\int_0^{\tau \exp(\theta_1 n)} f \circ h_t^+(x) dt = \sum_{i=1}^r \exp(n\theta_i) m_{\Phi_i^-}(f) \Phi_i^+(\sigma^n x, \tau) + O(n^{m+1}). \quad (52)$$

2.10 The Hölder property.

As above, we write $\Phi^+(x, t) = \Phi^+([x, h_t^+ x])$. Our next aim is to show that $\Phi^+(x, t)$ is Hölder in t for any $x \in X_\sigma$.

Proposition 15 *There exist positive constants C_Γ and t_0 , depending only on Γ such that the following is true. Let $v \in E^+$, $Qv = \exp(\theta)v$, $|v| = 1$. Then for all $x \in X$ and positive $t < t_0$ we have*

$$|\Phi_v^+(x, t)| \leq C_\Gamma t^{\Re \theta / \theta_1}.$$

Proposition 16 *There exist positive constants C_Γ and t_0 , depending only on Γ such that the following is true. Let $v_0 = 0$, $v_1, \dots, v_l \in E^+$, $Qv_i = \exp(\theta)v_i + v_{i-1}$. Assume $v \in \mathbb{C}v_1 \oplus \dots \oplus \mathbb{C}v_l$ satisfies $|v| = 1$. Then for all $x \in X$ and positive $t < t_0$ we have*

$$|\Phi_v^+(x, t)| \leq C_\Gamma |\log t|^{l-1} t^{\Re \theta / \theta_1}.$$

Proof of Propositions 15, 16. Denote $\gamma = [x, h_t^+ x]$. If t is small enough, then $\hat{\gamma}(0) = \emptyset$. Let n_0 be the smallest positive integer such that $\hat{\gamma}(n_0) \neq \emptyset$. There exist positive constants C_1, C_2 , depending only on Γ , such that

$$C_1 t \leq \exp(-\theta_1 n_0) \leq C_2 t,$$

and Propositions 15, 16 follow now from Proposition 13.

Corollary 2 *There exist positive constants $\theta > 0$ and $t_0 > 0$ depending only on Q such that for all $v \in E^+$, $|v| = 1$, all $x \in X$ and all positive $t < t_0$ we have*

$$|\Phi_v^+(x, t)| \leq t^{\theta/\theta_1}.$$

For $v \in E^+$, $|v| = 1$ denote

$$\theta_v = \lim_{n \rightarrow \infty} \frac{\log |Q^n v|}{n}.$$

Corollary 3 *For any $\varepsilon > 0$ there exists a constant T_ε depending only on ε and Γ such that for any $v \in E^+$, $|v| = 1$, any $x \in X$ and any $T > T_\varepsilon$, we have*

$$|\Phi_v^+(x, T)| \leq T^{\theta_v/\theta_1 + \varepsilon}.$$

Proof: Indeed, let t_0 be the constant given by Proposition 16. Let $n_0 = n_0(T)$ be the smallest such integer that $T = \tau \exp(n(T)\theta_1)$, where $\tau < t_0$. Since $\Phi_v^+(x, T) = \Phi_{Q^{n_0}v}^+(\sigma^{n_0}x, \tau)$ for all n , it follows from Proposition 16 that

$$|\Phi^+(x, T)| \leq C_\Gamma n_0^{m+1} \exp(n_0 \Re(\theta_v)) \leq C_\Gamma T^{\theta_v/\theta_1 + \varepsilon}$$

if T is sufficiently large (depending only on ε).

Corollary 4 *For any $v \in E^+$ we have*

$$\limsup_{T \rightarrow \infty} \frac{\log |\Phi_v^+(x, T)|}{\log T} = \frac{\theta_v}{\theta_1}. \quad (53)$$

Indeed, the upper bound for the limit superior follows from Corollary 3, and the lower bound is immediate from the relation $\Phi_v^+(\gamma_n(x)) = (Q^n v)_{F(x_{n+1})}$.

Corollary 5 *For any $\tau \in \mathbb{R}$ and any $v \in E^+$ satisfying $v \neq 0$, $\sum_{i=1}^m v_i \lambda_i = 0$, the function $\Phi_v^+(x, \tau)$ is not a constant in x .*

Proof: Indeed, assume $\Phi_v^+(x, \tau) = c$ identically. Then $\Phi^+(x, k\tau) = kc$, which contradicts (53): if $c = 0$, then the limit superior is 0; if $c \neq 0$, then the limit superior is 1.

2.11 Tightness.

In this subsection, we assume that Q has a simple real second eigenvalue $\exp(\theta_2)$, $\theta_2 > 0$. Let v_2 be the corresponding eigenvector and let $\Phi_2^+ = \mathcal{I}(v_2)$. Take $x \in X$ and consider $\Phi^+(x, \tau)$ as a continuous function of τ on the unit interval. Let η be the distribution of $\Phi_2^+(x, \tau)$ in $C[0, 1]$. Note that by Corollary 5, for any τ_0 the value of $\Phi_2^+(x, \tau)$ is not constant on X , so the measure η is nondegenerate.

Let $\mathfrak{S}_n[f, x]$ be defined by the equation (47). Introduce a sequence of measures μ_n on $C[0, 1]$ by the formula $\mu_n = \mathfrak{S}[n, f]_* \nu_\Gamma$.

By Theorem 8.1 in Billingsley [3], p.54, to prove Theorem 6 it suffices to establish the following two Lemmas.

Lemma 5 *Finite-dimensional distributions of the measures μ_n weakly converge to those of η .*

Lemma 6 *The family μ_n is tight in $C[0, 1]$.*

Proof of Lemma 5. By Theorem 5

$$\int_0^T f \circ h_t^+(x) dt = \Phi_f^+(x, T) + O((\log T)^{m+1}).$$

Let v_2 be the eigenvector corresponding to the eigenvalue $\exp(\theta_2)$, $|v| = 1$, and let $\Phi_2^+ \in \mathcal{Y}^+$ be the corresponding measure. We have

$$E^+ = \mathbb{C}v_2 \oplus E_3,$$

where E_3 is spanned by Jordan cells corresponding to eigenvalues with absolute value less than $\exp(\theta_2)$. Let ζ be a number smaller than θ_2 but greater than the spectral radius of $Q|_{E_3}$. Write

$$\Phi_f^+ = \alpha(f)\Phi_2^+ + \beta(f)\Phi_{v_3}^+, \quad (54)$$

where $v_3 \in E^+$, $|v_3| = 1$, and $\alpha(f), \beta(f)$ are continuous functionals on $Lip_w^+(X)$, so, in particular, we have

$$|\alpha(f)| < C_{01}\|f\|_{Lip_w^+}; \quad |\beta(f)| < C_{02}\|f\|_{Lip_w^+},$$

where the constants C_{01}, C_{02} only depend on Γ .

By Corollary 2, there exists t_0 depending only on Γ such that for any positive t such that $t < t_0$, any $x \in X$ and any $v \in E^+$ satisfying $|v| = 1$ we have

$$|\Phi_v^+(x, t)| \leq 1. \quad (55)$$

Write $T = t \exp(n\theta_1)$, where $t < t_0$. Since $\Phi_{v_3}^+(x, T) = \Phi_{Q^n v_3}^+(\sigma^n x, t)$, for all sufficiently large n , we have $|Q^n v_3| < \exp(\zeta n)$ and therefore

$$|\Phi_{v_3}^+(x, \tau \exp(n\theta_1))| < \exp(n\zeta) \quad (56)$$

for all $x \in X$. By Theorem 5 we have

$$\left| \int_0^{\tau \exp(n\theta_1)} f \circ h_t^+(x) dt - \Phi_f^+(x, \tau \exp(n\theta_1)) \right| = O(n^{m+1}). \quad (57)$$

Since

$$\Phi_f^+(x, \tau \exp(n\theta_1)) = \alpha(f)\Phi_2^+((x, \tau \exp(n\theta_1))) + \beta(f)\Phi_{v_3}^+(x, \tau \exp(n\theta_1))$$

combining the equality

$$\Phi_2^+(x, \tau \exp(n\theta_1)) = \exp(n\theta_2)\Phi_2^+(\sigma^n x, \tau)$$

with the bound (56), we obtain, for all large n and all $x \in X$, uniformly in $\tau \in [0, 1]$, the estimate

$$|\mathfrak{S}_n[f, x](\tau) - \alpha(f)\Phi_2^+(\sigma^n x, \tau)| \leq C_\Gamma \|f\|_{Lip_w^+} \exp((\zeta - \theta_2)n).$$

Since σ preserves the measure ν , it follows that the k -dimensional distributions of $(\mathfrak{S}_n[f, x](\tau_1), \mathfrak{S}_n[f, x](\tau_2), \dots, \mathfrak{S}_n[f, x](\tau_k))$ converge to the k -dimensional distribution of $(\Phi_2^+(x, \tau_1), \Phi_2^+(x, \tau_2), \dots, \Phi_2^+(x, \tau_k))$, and Lemma 5 is proved.

The argument above yields also

Proposition 17 *There exist positive constants $C_0 = C_0(\Gamma)$ and $T_0 = T_0(\Gamma)$ such that for any $x \in X$, any $f \in Lip_{w,0}^+(X)$ and any $T > T_0$ we have*

$$|\int_0^T f \circ h_t^+(x) dt| \leq C_0 \cdot \|f\|_{Lip_w^+} \cdot T^{\theta_2/\theta_1}.$$

Indeed, for sufficiently large T , $T = t \exp(n\theta_1)$, where $t < t_0$, from (54) we have

$$\Phi_f^+(x, T) = \alpha(f) \exp(n\theta_2) \Phi_2^+(\sigma^n x, t) + O(\exp(n\zeta)).$$

Since, by (55), we have $|\Phi_2^+(\sigma^n x, t)| \leq 1$, Proposition 17 is established.

We proceed to the proof of Lemma 6.

Proposition 18 *There exists a constant C_Γ depending only on Γ such that for any $f \in Lip_{w,0}^+(X)$, any $n > 0$, any $x \in X$ and any $\tau_1, \tau_2 \in [0, 1]$, we have*

$$|\mathfrak{S}_n[x, f](\tau_2) - \mathfrak{S}_n[x, f](\tau_1)| \leq C_\Gamma \|f\|_{Lip_w^+} |\tau_2 - \tau_1|^{\theta_2/\theta_1}.$$

Lemma 6 follows from Proposition 18 by the Arzelà-Ascoli Theorem.

Proof of Proposition 18: Let $\tau_1, \tau_2 \in [0, 1]$, $\tau_1 < \tau_2$. For brevity, write $\mathfrak{S}_n = \mathfrak{S}_n[f, x]$. We have then

$$\mathfrak{S}_n(\tau_2) - \mathfrak{S}_n(\tau_1) = \frac{1}{\exp(n\theta_2)} \int_{\tau_1 \exp(n\theta_1)}^{\tau_2 \exp(n\theta_1)} f \circ h_t^+(x) dt.$$

Let T_0 be the constant given by Proposition 17 and assume first that

$$(\tau_2 - \tau_1) \cdot \exp(n\theta_1) \geq T_0.$$

By Proposition 17 we have

$$\int_{\tau_1 \exp(n\theta_1)}^{\tau_2 \exp(n\theta_1)} f \circ h_t(x) dt \leq C \|f\|_{Lip_w^+} \cdot (\tau_2 - \tau_1)^{\theta_2/\theta_1} \exp(n\theta_2),$$

and, consequently,

$$|\mathfrak{S}_n(\tau_2) - \mathfrak{S}_n(\tau_1)| \leq C_{33} (\tau_2 - \tau_1)^{\theta_2/\theta_1},$$

where the constant C_{33} only depends on Γ .

Now let $\tau_2 - \tau_1 = \tau_0 \exp(-n\theta_1)$, $\tau_0 < T_0$. Since

$$\exp(-n\theta_2) = ((\tau_2 - \tau_1)/\tau_0)^{\theta_2/\theta_1},$$

using boundedness of f , write

$$\begin{aligned} \frac{1}{\exp(n\theta_2)} \int_{\tau_1 \exp(n\theta_1)}^{\tau_2 \exp(n\theta_1)} f \circ h_t^+(x) dt &\leq \exp(-n\theta_2) \cdot \|f\|_\infty \cdot \tau_0 \leq \\ &\leq \tau_0^{1-\theta_2/\theta_1} \|f\|_\infty (\tau_2 - \tau_1)^{\theta_2/\theta_1} \leq T_0^{1-\theta_2/\theta_1} \|f\|_\infty (\tau_2 - \tau_1)^{\theta_2/\theta_1}, \end{aligned}$$

and the Proposition is proved. Theorem 6 is proved completely.

2.12 A symbolic coding for translation flows on surfaces.

To derive Theorems 1, 2 from Theorems 5, 6, it remains to observe that the vertical flow on the stable foliation of a pseudo-Anosov diffeomorphism is isomorphic to a symbolic flow on the asymptotic foliation of a Markov compactum obtained from the decomposition of the underlying surface into Veech's zippered rectangles, see [4], Sec. 4. The identification of E^+ (and, consequently, of \mathcal{Y}^+) with the corresponding subspace in cohomology is given by Proposition 4.16 in Veech[14]. The fact that the pairing between cocycles corresponds to the cup-product is immediate from Proposition 4.19 in [14].

3 Spaces of Markov Compacta.

Let \mathfrak{G} be the set of all oriented graphs on m vertices such that there is an edge starting at every vertex and an edge ending at every vertex. As before, for a graph $\Gamma \in \mathfrak{G}$, we denote by $\mathcal{E}(\Gamma)$ the set of its edges and by $A(\Gamma)$ its incidence matrix: $A_{ij}(\Gamma) = \#\{e \in \mathcal{E}(\Gamma) : I(e) = i, F(e) = j\}$. Denote $\Omega = \mathfrak{G}^{\mathbb{Z}}$:

$$\Omega = \{\omega = \dots \omega_{-n} \dots \omega_n \dots, \omega_i \in \mathfrak{G}, i \in \mathbb{Z}\},$$

For $\omega \in \Omega$, denote by $X(\omega)$ the corresponding Markov compactum:

$$X(\omega) = \{x = \dots x_{-n} \dots x_n \dots, x_n \in \mathcal{E}(\omega_n), F(x_{n+1}) = I(x_n)\}.$$

For $x \in X$, $n \in \mathbb{Z}$, introduce the sets

$$\gamma_n^+(x) = \{x' \in X(\omega) : x'_t = x_t, t \geq n\}; \quad \gamma_n^-(x) = \{x' \in X(\omega) : x'_t = x_t, t \leq n\};$$

$$\gamma_\infty^+(x) = \bigcup_{n \in \mathbb{Z}} \gamma_n^+(x); \quad \gamma_\infty^-(x) = \bigcup_{n \in \mathbb{Z}} \gamma_n^-(x).$$

The sets $\gamma_\infty^+(x)$ are leaves of the asymptotic foliation \mathcal{F}_ω^+ on $X(\omega)$; the sets $\gamma_\infty^-(x)$ are leaves of the asymptotic foliation \mathcal{F}_ω^- on $X(\omega)$.

For $n \in \mathbb{Z}$ let $\mathfrak{C}_{n,\omega}^+$ be the collection of all subsets of $X(\omega)$ of the form $\gamma_n^+(x)$, $n \in \mathbb{Z}$, $x \in X$; similarly, $\mathfrak{C}_{n,\omega}^-$ is the collection of all subsets of the form $\gamma_n^-(x)$. Set

$$\mathfrak{C}_\omega^+ = \bigcup_{n \in \mathbb{Z}} \mathfrak{C}_{n,\omega}^+; \mathfrak{C}_\omega^- = \bigcup_{n \in \mathbb{Z}} \mathfrak{C}_{n,\omega}^-. \quad (58)$$

Just as in the periodic case, the collections $\mathfrak{C}_{n,\omega}^+$, $\mathfrak{C}_{n,\omega}^-$, \mathfrak{C}_ω^+ , \mathfrak{C}_ω^- are semi-rings.

Remark. To make notation lighter, we shall often omit the subscript ω and only include it when dependence on ω is underlined.

3.1 Measures and Cocycles.

Let σ be the shift on Ω given by the formula $(\sigma\omega)_n = \omega_{n+1}$. Let \mathbb{P} be an ergodic σ -invariant probability measure on Ω . We then have a natural cocycle \mathbb{A} on the system $(\Omega, \sigma, \mathbb{P})$ defined, for $n > 0$, by the formula

$$\mathbb{A}(n, \omega) = A(\omega_n) \dots A(\omega_1).$$

The cocycle \mathbb{A} will be called the *renormalization cocycle*.

We need the following assumptions on the measure \mathbb{P} and on the cocycle \mathbb{A} .

Assumption 1 *The matrices $A(\omega_n)$ are almost surely invertible with respect to \mathbb{P} . There exists $\Gamma \in \mathfrak{G}$ such that $\mathbb{P}(\Gamma) > 0$.*

Assumption 2 *The logarithm of the renormalization cocycle (and of its inverse) is integrable.*

For $n < 0$ set

$$\mathbb{A}(n, \omega) = A^{-1}(\omega_{-n}) \dots A^{-1}(\omega_0).$$

and set $\mathbb{A}(0, \omega)$ to be the identity matrix.

The *transpose* cocycle \mathbb{A}^t over the dynamical system $(\Omega, \sigma^{-1}, \mathbb{P})$ defined, for $n > 0$, by the formula

$$\mathbb{A}^t(n, \omega) = A^t(\omega_{1-n}) \dots A^t(\omega_0).$$

Similarly, for $n < 0$ write

$$\mathbb{A}^t(n, \omega) = (A^t)^{-1}(\omega_{-n}) \dots (A^t)^{-1}(\omega_1).$$

and set $\mathbb{A}^t(0, \omega)$ to be the identity matrix.

By Assumptions 1, 2, for \mathbb{P} -almost any $\omega \in \Omega$ we have the decompositions

$$\mathbb{R}^m = E_\omega^+ \oplus E_\omega^-; \mathbb{R}^m = \tilde{E}_\omega^+ \oplus \tilde{E}_\omega^-,$$

where E^+ is the Lyapunov subspace corresponding to positive Lyapunov exponents of \mathbb{A} ; \tilde{E}^+ is the Lyapunov subspace corresponding to positive Lyapunov exponents of \mathbb{A}^t ; E^- is the Lyapunov subspace corresponding to zero and negative Lyapunov exponents of \mathbb{A} ; \tilde{E}^- is the Lyapunov subspace corresponding to

zero and negative Lyapunov exponents of \mathbb{A}^t . The standard inner product on \mathbb{R}^m yields a nondegenerate pairing between the spaces E_ω^+ and \tilde{E}_ω^+ .

In particular, by Assumption 1, the spaces E_ω^+ and \tilde{E}_ω^+ each contain a unique vector all whose coordinates are positive; we denote these vectors by $h^{(\omega)}$ and $\lambda^{(\omega)}$, respectively, and assume that they are normalized by (12).

Let $v \in E_\omega^+$ and for all $n \in \mathbb{Z}$ set $v^{(n)} = \mathbb{A}(n, \omega)v$. Introduce a finitely-additive complex-valued measure Φ_v^+ on the semi-ring \mathfrak{C}_ω^+ (defined in (58)) by the formula

$$\Phi_v^+(\gamma_{n+1}^+(x)) = (v^{(n)})_{F(x_{n+1})}. \quad (59)$$

As before, the measure Φ_v^+ is invariant under holonomy along \mathcal{F}^- : by definition, we have the following

Proposition 19 *If $F(x_n) = F(x'_n)$, then $\Phi_v^+(\gamma_n^+(x)) = \Phi_v^+(\gamma_n^+(x'))$.*

The measures Φ_v^+ span a complex linear space, which is denoted \mathcal{Y}_ω^+ . The map $\mathcal{I}_\omega : v \rightarrow \Phi_v^+$ is an isomorphism between E_ω^+ and \mathcal{Y}_ω^+ . Set $\Phi_{1,\omega}^+ = \mathcal{I}_\omega(h^{(\omega)})$.

Now for $\tilde{v} \in \tilde{E}^+$ and for all $n \in \mathbb{Z}$ set $\tilde{v}^{(n)} = \mathbb{A}^t(n, \omega)\tilde{v}$ and introduce a finitely-additive complex-valued measure $\Phi_{\tilde{v}}^-$ on the semi-ring \mathfrak{C}_ω^- (defined in (58)) by the formula

$$\Phi_{\tilde{v}}^-(\gamma_n^-(x)) = (\tilde{v}^{(-n)})_{I(x_n)}. \quad (60)$$

By definition, the measure $\Phi_{\tilde{v}}^-$ is invariant under holonomy along \mathcal{F}^+ : more precisely, we have the following

Proposition 20 *If $I(x_n) = I(x'_n)$, then $\Phi_{\tilde{v}}^-(\gamma_n^-(x)) = \Phi_{\tilde{v}}^-(\gamma_n^-(x'))$.*

Let \mathcal{Y}_ω^- be the space spanned by the measures $\Phi_{\tilde{v}}^-$, $\tilde{v} \in \tilde{E}^+$. The map $\tilde{\mathcal{I}}_\omega : \tilde{v} \rightarrow \Phi_{\tilde{v}}^-$ is an isomorphism between \tilde{E}_ω^+ and \mathcal{Y}_ω^- . Set $\Phi_{1,\omega}^- = \tilde{\mathcal{I}}_\omega(\lambda^{(\omega)})$.

Define a map $t_\sigma : X_\omega \rightarrow X_{\sigma\omega}$ by $(t_\sigma x)_i = x_{i+1}$. The map t_σ induces a map $t_\sigma^* : \mathcal{Y}_{\sigma\omega}^+ \rightarrow \mathcal{Y}_\omega^+$ given, for $\Phi_{\sigma\omega}^+ \in \mathcal{Y}_{\sigma\omega}^+$ and $\gamma \in \mathfrak{C}_\omega^+$, by the formula

$$t_\sigma^* \Phi_{\sigma\omega}^+(\gamma) = \Phi_{\sigma\omega}^+(t_\sigma \gamma).$$

We have the following commutative diagrams:

$$\begin{array}{ccc} E_\omega^+ & \xrightarrow{\mathcal{I}_\omega} & \mathcal{Y}_\omega^+ \\ \downarrow \mathbb{A}(1, \omega) & & \uparrow t_\sigma^* \\ E_{\sigma\omega}^+ & \xrightarrow{\mathcal{I}_{\sigma\omega}} & \mathcal{Y}_{\sigma\omega}^+ \\ \\ \tilde{E}_\omega^+ & \xrightarrow{\tilde{\mathcal{I}}_\omega} & \mathcal{Y}_\omega^- \\ \uparrow \mathbb{A}^t(1, \sigma\omega) & & \uparrow t_\sigma^* \\ \tilde{E}_{\sigma\omega}^+ & \xrightarrow{\tilde{\mathcal{I}}_{\sigma\omega}} & \mathcal{Y}_{\sigma\omega}^- \end{array}$$

3.2 Pairings and weakly Lipschitz functions.

Given $\Phi^+ \in \mathcal{Y}_\omega^+$, $\Phi^- \in \mathcal{Y}_\omega^-$, introduce a finitely additive measure $\Phi^+ \times \Phi^-$ on the semi-ring \mathfrak{C} of cylinders in $X(\omega)$: for any $C \in \mathfrak{C}$ and $x \in C$, set

$$\Phi^+ \times \Phi^-(C) = \Phi^+(\gamma_\infty^+(x) \cap C) \cdot \Phi^-(\gamma_\infty^-(x) \cap C). \quad (61)$$

Note that by Propositions 19, 20, the right-hand side in (61) does not depend on $x \in C$.

As above, for $\Phi^- \in \mathcal{Y}_\omega^-$, denote

$$m_{\Phi^-} = \Phi_1^+ \times \Phi^-. \quad (62)$$

In particular, we have a positive countably additive measure

$$\nu_\omega = \Phi_{h(\omega)}^+ \times \Phi_{\lambda(\omega)}^-.$$

There is a natural \mathbb{C} -linear pairing \langle, \rangle between the spaces \mathcal{Y}_ω^+ and \mathcal{Y}_ω^- : for $\Phi^+ \in \mathcal{Y}_\omega^+$, $\Phi^- \in \mathcal{Y}_\omega^-$, set

$$\langle \Phi^+, \Phi^- \rangle = \Phi^+ \times \Phi^-(X(\omega)). \quad (63)$$

As in Sec. 2.3, we have

Proposition 21 *Let $v \in E_\omega^+$, $\tilde{v} \in \tilde{E}_\omega^+$, $\Phi_v^+ = \mathcal{I}_\omega(v)$, $\Phi_{\tilde{v}}^- = \tilde{\mathcal{I}}_\omega(\tilde{v})$. Then*

$$\langle \Phi_v^+, \Phi_{\tilde{v}}^- \rangle = \sum_{i=1}^m v_i \tilde{v}_i. \quad (64)$$

The pairing \langle, \rangle is non-degenerate and t_σ^ -invariant.*

The function space $Lip_w^+(X(\omega))$ is introduced in the same way as before: a bounded Borel-measurable function $f : X(\omega) \rightarrow \mathbb{C}$ belongs to the space $Lip_w^+(X)$ if there exists a constant $C > 0$ such that for all $n \geq 0$ and any $x, x' \in X$ satisfying $F(x_{n+1}) = F(x'_{n+1})$, we have

$$\left| \int_{\gamma_n^+(x)} f d\Phi_1^+ - \int_{\gamma_n^+(x')} f d\Phi_1^+ \right| \leq C, \quad (65)$$

and, if C_f is the infimum of all C satisfying (65), then we norm $Lip_w^+(X)$ by setting

$$\|f\|_{Lip_w^+} = \sup_X f + C_f.$$

As before, we denote by $Lip_{w,0}^+(X(\omega))$ the subspace of functions of ν_ω -integral zero.

Take $\Phi^- \in \mathcal{Y}^-$. Any function $f \in Lip_w^+(X)$ is integrable with respect to the measure m_{Φ^-} in the same sense as in Sec. 2.4, and a measure $\Phi_f^+ \in \mathcal{Y}^+$ is defined by the requirement that for any $\Phi^- \in \mathcal{Y}^-$ we have

$$\langle \Phi_f^+, \Phi^- \rangle = \int_{X(\omega)} f dm_{\Phi^-}. \quad (66)$$

Note that the mapping $\Xi_\omega^+ : Lip_w^+(X(\omega)) \rightarrow \mathcal{Y}_\omega^+$ given by $\Xi_\omega^+(f) = \Phi_f^+$ is continuous by definition and satisfies

$$\Xi_{\sigma\omega}^+(f \circ t_\sigma) = (t_\sigma)^* \Xi_\omega^+(f). \quad (67)$$

From the definitions we also have

Proposition 22 *Let $\Phi^+(1), \dots, \Phi^+(r)$ be a basis in \mathcal{Y}_ω^+ and let $\Phi^-(1), \dots, \Phi^-(r)$ be the dual basis in \mathcal{Y}_ω^- with respect to the pairing \langle, \rangle . Then for any $f \in Lip_w^+(X(\omega))$ we have*

$$\Phi_f^+ = \sum_{i=1}^r (m_{\Phi^-(i)}(f)) \Phi^+(i).$$

3.3 Orderings and flows.

Assume that for \mathbb{P} -almost every ω a partial ordering $\mathfrak{o}(\omega)$ is given on $\mathcal{E}(\omega_n)$ for all $n \in \mathbb{Z}$ in such a way that edges starting at a given vertex are ordered linearly, while edges starting at different vertices are incomparable. Assume, moreover, that the orders $\mathfrak{o}(\omega)$ are σ -invariant, in the sense that the ordering $\mathfrak{o}(\omega)$ on $\mathcal{E}(\omega_n)$ is the same as the ordering $\mathfrak{o}(\sigma\omega)$ on $\mathcal{E}((\sigma\omega)_{n-1})$.

Similarly to the above, construct spaces $X_\mathfrak{o}(\omega)$ and introduce a flow $h_t^{(+,\omega)}$ on each $X_\mathfrak{o}(\omega)$. The shift σ renormalizes the flows $h_t^{(+,\omega)}$: if we set

$$H^{(1)}(n, \omega) = \|\mathbb{A}(n, \omega)\|, \quad (68)$$

then for any $t \in \mathbb{R}$ we have a commutative diagram

$$\begin{array}{ccc} X(\omega) & \xrightarrow{h_t^{(+,\omega)}} & X(\omega) \\ \downarrow t_\sigma & & \downarrow t_\sigma \\ X(\sigma\omega) & \xrightarrow{h_{t/H^{(1)}(1,\omega)}^{(+,\sigma\omega)}} & X(\sigma\omega) \end{array}$$

As before, each measure $\Phi^+ \in \mathcal{Y}_\omega^+$ yields a Hölder cocycle over the flow $h_t^{(+,\omega)}$; we shall denote the cocycle by the same letter as the measure.

Note that for any $\Phi^- \in \mathcal{Y}_\omega^-$ the measure m_{Φ^-} defined by (62) satisfies

$$(h_t^{(+,\omega)})_* m_{\Phi^-} = m_{\Phi^-},$$

similarly to G. Forni's invariant distributions [5], [6].

Note that the mapping $\Xi_\omega^+ : Lip_w^+(X(\omega)) \rightarrow \mathcal{Y}_\omega^+$ given by $\Xi_\omega^+(f) = \Phi_f^+$ by definition satisfies

$$\Xi_\omega^+(f \circ h_t^{(+,\omega)}) = \Xi_\omega^+(f). \quad (69)$$

We thus have the following

Theorem 7 Let \mathbb{P} be an ergodic σ -invariant probability measure on Ω satisfying the assumptions 1, 2. For any $\varepsilon > 0$ there exists a positive constant C_ε depending only on \mathbb{P} such that the following holds. For \mathbb{P} -almost any ω there exists a continuous mapping $\Xi_\omega^+ : Lip_w^+(X(\omega)) \rightarrow \mathcal{Y}_\omega^+$ such that for any $f \in Lip_w^+(X(\omega))$, any $x \in X(\omega)$ and all $T > 0$ we have

$$\left| \int_0^T f \circ h_t^{(+, \omega)}(x) dt - \Xi_\omega^+(f)(x, t) \right| \leq C_\varepsilon \|f\|_{Lip_w^+} (1 + T^\varepsilon).$$

The mapping Ξ_ω^+ satisfies the equality $\Xi_\omega^+(f \circ h_t^{(+, \omega)}) = \Xi_\omega^+(f)$. The diagram

$$\begin{array}{ccc} Lip_w^+(X(\sigma\omega)) & \xrightarrow{\Xi_{\sigma\omega}^+} & \mathcal{Y}_{\sigma\omega}^+ \\ \downarrow t_\sigma^* & & \downarrow t_\sigma^* \\ Lip_w^+(X(\omega)) & \xrightarrow{\Xi_\omega^+} & \mathcal{Y}_\omega^+ \end{array}$$

is commutative.

The mapping Ξ_ω^+ is given by $\Xi_\omega^+(f) = \Phi_f^+$, where Φ_f^+ is defined by (66).

Now assume that the second Lyapunov exponent θ_2 of the renormalization cocycle \mathbb{A} is positive and simple. Let $v_2 \in E_\omega^+$ be a Lyapunov vector corresponding to the exponent $\exp(\theta_2)$ (such a vector is defined up to multiplication by a scalar). Introduce a multiplicative cocycle $H^{(2)}(n, \omega)$ over σ by the formula

$$H^{(2)}(n, \omega) = \frac{|\mathbb{A}(n, \omega)v_2^{(\omega)}|}{|v_2^{(\omega)}|}. \quad (70)$$

Recall that the cocycle $H^{(1)}(n, \omega)$ is given by (68). Similarly to the above, given a bounded measurable function $f : X(\omega) \rightarrow \mathbb{R}$ and $x \in X(\omega)$, introduce a continuous function $\mathfrak{S}_n[f, x]$ on the unit interval by the formula

$$\mathfrak{S}_n[f, x](\tau) = \int_0^{\tau H^{(1)}(n, \omega)} f \circ h_t^{(+, \omega)}(x) dt. \quad (71)$$

The functions $\mathfrak{S}_n[f, x]$ are $C[0, 1]$ -valued random variables on the probability space $(X(\omega), \nu_\omega)$.

Theorem 8 Let \mathbb{P} be an ergodic σ -invariant probability measure on Ω satisfying the assumptions 1, 2 and such the second Lyapunov exponent of the renormalization cocycle \mathbb{A} with respect to \mathbb{P} is positive and simple.

For \mathbb{P} -almost any $\omega' \in \Omega$ there exists a non-degenerate compactly supported measure $\eta_{\omega'}$ on $C[0, 1]$ and, for \mathbb{P} -almost any pair (ω, ω') there exists a sequence of moments $l_n = l_n(\omega, \omega')$ such that the following holds.

For \mathbb{P} -almost any ω there exists a continuous functional

$$\mathfrak{a}^{(\omega)} : Lip_w^+(X(\omega)) \rightarrow \mathbb{R}$$

such that for \mathbb{P} -almost any ω' and any $f \in Lip_{w,0}^+(X(\omega))$ satisfying $\mathfrak{a}^{(\omega)}(f) \neq 0$ the sequence of random variables

$$\frac{\mathfrak{S}_{l_n(\omega, \omega')}[f, x]}{\mathfrak{a}^{(\omega)}(f) H^{(2)}(l_n(\omega, \omega'), \omega)}$$

converges in distribution to $\eta_{\omega'}$ as $n \rightarrow \infty$.

Theorems 7, 8 imply Theorems 3, 4. The proofs of Theorems 7, 8 follow the same pattern as those of Theorems 5, 6; detailed proofs will appear in the sequel to this paper.

References

- [1] Adamczewski, Boris. Symbolic discrepancy and self-similar dynamics. Ann. Inst. Fourier (Grenoble) 54 (2004), no. 7, 2201–2234.
- [2] Avila, Artur; Viana, Marcelo. Simplicity of Lyapunov spectra: proof of the Zorich-Kontsevich conjecture. Acta Math. 198 (2007), no. 1, 1–56.
- [3] Billingsley, Patrick. Convergence of probability measures. Wiley, 1968.
- [4] Bufetov, A. Suspension flows over Vershik’s automorphisms, preprint, www.arxiv.org.
- [5] Forni, Giovanni. Solutions of the cohomological equation for area-preserving flows on compact surfaces of higher genus. Ann. of Math. (2) 146 (1997), no. 2, 295–344.
- [6] Forni, Giovanni. Deviation of ergodic averages for area-preserving flows on surfaces of higher genus. Ann. of Math. (2) 155 (2002).
- [7] Ito, Shunji. A construction of transversal flows for maximal Markov automorphisms. Tokyo J. Math. 1 (1978), no. 2, 305–324.
- [8] M.Kontsevich, Lyapunov exponents and Hodge theory, “Mathematical Beauty of Physics”, Saclay, 1996.
- [9] Marmi, S.; Moussa, P.; Yoccoz, J.-C. The cohomological equation for Roth-type interval exchange maps. J. Amer. Math. Soc. 18 (2005), no. 4, 823–872.
- [10] H. Masur, Interval exchange transformations and measured foliations. Ann. of Math. (2) 115 (1982), no. 1, 169–200.
- [11] Sinai, Ya. G., Ulcigrai, C. Weak mixing in interval exchange transformations of periodic type. Lett. Math. Phys. 74 (2005), no. 2, 111–133.
- [12] Veech, William A. Gauss measures for transformations on the space of interval exchange maps. Ann. of Math. (2) 115 (1982), no. 1, 201–242.

- [13] Veech, William A. The Teichmüller geodesic flow. *Ann. of Math.* (2) 124 (1986), no. 3, 441–530.
- [14] Veech, William A. Moduli spaces of quadratic differentials. *J. Analyse Math.* 55 (1990), 117–171.
- [15] Vershik, A. M. A theorem on Markov periodic approximation in ergodic theory. (Russian) Boundary value problems of mathematical physics and related questions in the theory of functions, 14. *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* 115 (1982), 72–82, 306.
- [16] Vershik, A. M. The adic realizations of the ergodic actions with the homeomorphisms of the Markov compact and the ordered Bratteli diagrams. *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* 223 (1995), Teor. Predstav. Din. Sistemy, Kombin. i Algoritm. Metody. I, 120–126, 338; translation in *J. Math. Sci. (New York)* 87 (1997), no. 6, 4054–4058.
- [17] Vershik, A. M.; Livshits, A. N. Adic models of ergodic transformations, spectral theory, substitutions, and related topics. Representation theory and dynamical systems, 185–204, *Adv. Soviet Math.*, 9, Amer. Math. Soc., Providence, RI, 1992.
- [18] Zorich, Anton. Deviation for interval exchange transformations. *Ergodic Theory Dynam. Systems* 17 (1997), no. 6, 1477–1499.